

1   **Title:** Caffeinated soda intake in children is associated with neurocognitive vulnerabilities of  
2   substance misuse and is predictive of a higher risk of alcohol sipping after 12 months

3

4   **Authors:**

5   Mina Kwon<sup>1</sup>, Hyeonjin Kim<sup>1</sup>, Jaeyeong Yang<sup>1</sup>, Yoseph Lee<sup>1</sup>, Jihyun K. Hur<sup>1,2</sup>, Tae-Ho Lee<sup>3</sup>, and  
6   Woo-Young Ahn<sup>1,4</sup>

7

8   <sup>1</sup>Department of Psychology, Seoul National University, Seoul, Korea 08826

9   <sup>2</sup>Department of Psychology, Yale University, New Haven, CT 06511

10   <sup>3</sup>Department of Psychology, Virginia Tech, Blacksburg, VA 24060

11   <sup>4</sup>Department of Brain and Cognitive Sciences, Seoul National University, Seoul, Korea 08826

12

13   **Corresponding author:**

14   Woo-Young Ahn, Ph.D.

15   Department of Psychology

16   Seoul National University

17   Seoul, Korea 08826

18   Tel: +82-2-880-2538, Fax: +82-2-877-6428. E-mail: [wahn55@snu.ac.kr](mailto:wahn55@snu.ac.kr)

19

20   **Running head:** Soda intake in children and alcohol sipping

21

22   **Word count:**

23   Number of Figures & Tables: 4 / 1

24   Number of words in the introduction: 662

25   Number of words in the method: 530

26   Number of words in the results: 779

27   Number of words in the discussion: 1,431

28

1   **Declaration of interests:** None to declare

2

3   **Funding:**

4   This work was supported by the Basic Science Research Program through the National Research  
5   Foundation (NRF) of Korea funded by the Ministry of Science, ICT, & Future Planning (NRF-  
6   2018R1C1B3007313 and NRF-2018R1A4A1025891); the Institute for Information &  
7   Communications Technology Planning & Evaluation (IITP) grant funded by the Korea government  
8   (MSIT) (No. 2019-0-01367, BabyMind); the National Research Foundation of Korea (NRF) grant  
9   funded by the Korea government (MSIT) (No. 2021M3E5D2A0102249311); and the Creative-  
10   Pioneering Researchers Program through Seoul National University to W.-Y.A.

## 1 ABSTRACT

2 **Background and Aims** Caffeinated soda contains two addictive substances, sugar and caffeine,  
3 and is the most preferred means of caffeine consumption among children. However, it remains  
4 unexplored if habitual caffeinated soda intake in childhood is associated with a higher risk of  
5 alcohol misuse in the future. Here, we investigated the neurocognitive correlates of caffeinated  
6 soda consumption and examined whether caffeinated soda intake is associated with a higher risk  
7 of alcohol initiation in children. **Methods** Using the Adolescent Brain Cognitive Development  
8 Study, we first investigated the relationships between frequent caffeinated soda intake and well-  
9 known risk factors of substance misuse: impaired working memory, high impulsivity, and blunted  
10 reward processing. We next examined whether caffeinated soda intake predicts more alcohol  
11 sipping after 12 months, and whether caffeinated soda intake mediates the link between  
12 neurocognitive risk factors and future alcohol sipping. **Results** Frequent consumption of  
13 caffeinated soda was associated with neurocognitive risk factors for substance misuse. Daily  
14 soda intake was associated with hyperactivity in the striatal and parietal regions and hypoactivity  
15 in the inferior frontal gyrus (IFG) during the emotional N-Back Task. In addition, the striatum, IFG,  
16 and anterior cingulate cortex (ACC) showed aberrant activity during the stop signal task, and the  
17 medial prefrontal cortex and the ACC exhibited increased and decreased activity, respectively,  
18 during the monetary incentive delay task. Furthermore, caffeinated soda intake predicted greater  
19 alcohol sipping after 12 months, and mediated the relationship between impulsivity and future  
20 alcohol sipping. **Conclusions** This study addresses the question that many parents are asking:  
21 "Is frequent caffeinated soda intake in childhood associated with a higher risk of substance  
22 misuse in the future?" We showed that caffeinated soda consumption in childhood was  
23 associated with impaired neurocognitive functioning and may predict future alcohol intake, which  
24 has significant public health implications.

25

## 1 INTRODUCTION

2 Soft drinks are commonly consumed even by children, and a vast majority of sodas  
3 contain caffeine [1–3]. Caffeinated soda refers to soft drinks containing caffeine as well as  
4 sweeteners and artificial flavorings. Thus, caffeinated soda combines the addictive substances of  
5 sugar and caffeine, both of which affect neurocognitive function and cause physical side effects,  
6 although the concept of sugar addiction remains controversial [4–6]. Not surprisingly, the  
7 consequences of excessive consumption of both sugar and caffeine have been well documented  
8 [7], including a strong association between caffeinated beverage consumption in adolescents and  
9 future substance use [8–13]. In prospective studies tracking the effects of substance use, the  
10 percentage of regular energy drink users who became alcohol or marijuana users after 1–2 years  
11 was approximately five times higher than that of non-energy drink users [9,12,13]. Others have  
12 shown that coffee or energy drink consumption in children significantly predicts future substance  
13 use, such as tobacco and alcohol use [8,10,12].

14 Previous studies examining the association between caffeine consumption and later  
15 substance use have focused more on adolescents who drink multiple caffeinated beverages on a  
16 daily basis, partially because the consumption level of coffee and energy drinks is extremely low  
17 in children (0.4% for coffee and < 0.1% for energy drinks in 9–10-year-old children) [14].  
18 However, caffeinated soda is the most preferred and accessible mode of caffeine intake in  
19 children [1,4,15]. Moreover, it is well known that adolescence is the most common period for  
20 initiating substance use, and an earlier onset of substance use predicts greater addiction severity  
21 [16–19]. Therefore, there is a critical need to investigate the association between caffeinated  
22 soda intake during preadolescent childhood and the risk of substance use disorders (SUDs) in  
23 the future, including vulnerabilities, such as associated neurocognitive deficits or early exposure  
24 to substances of abuse. Environmental factors, particularly the role of family, should be examined  
25 together, considering the previous studies that suggested a critical role of parental interventions  
26 in youth soft drink consumption and substance use [20–26]. Few studies have directly examined  
27 the effects of frequent caffeinated soda consumption in adolescents and children, and they either

1 targeted only behavioral aspects using surveys or self-reports without obtaining neural data, or  
2 targeted only adolescents and not children [9,27–29].

3 Here, we addressed the unanswered question of whether frequent consumption of  
4 caffeinated soda in children indicates a higher risk of later substance misuse, using the  
5 Adolescent Brain Cognitive Development (ABCD) Study. First, we examined the relationship  
6 between caffeinated soda intake and three well-known neurocognitive risk factors for SUDs in  
7 children: impaired working memory (WM), high impulsivity, and altered reward processing [30].  
8 These are the three main risk factors targeted in the ABCD study due to their significant  
9 implications for addiction (see [30] for more detail). We used psychological, behavioral, and  
10 neurological indices related to each risk factor, from the baseline data of the ABCD study (**Figure**  
11 **1**). We applied a machine learning approach to the indices so that we could identify multivariate  
12 risk factors for SUDs associated with caffeinated soda intake [31]. We next examined whether  
13 caffeinated soda intake could predict future alcohol sipping, as (1) alcohol sipping has been  
14 reported to predict future alcohol abuse [32–34], and (2) alcohol sipping is the most common  
15 gateway behavior toward other substances of abuse [19,35]. Lastly, we examined mediation  
16 models explaining how the interaction between daily caffeinated soda intake and the SUD risk  
17 factors leads to future alcohol sipping behavior. Based on the previous literature suggesting that  
18 caffeinated beverage consumption predicts future substance use [8–10,12,13], we hypothesized  
19 that caffeinated soda intake would mediate the relationship between neurocognitive risk factors  
20 and future alcohol sipping (**Figure 1A**). We also evaluated family risk factors as mediators of this  
21 relationship, based on the literature that family risk factors, including family substance use or  
22 family support, are risk factors for substance use in adolescents [24–26]. Taken together, we  
23 aimed to elucidate the potential underlying risks of frequent caffeinated soda intake during  
24 childhood.

25  
26

1    **METHODS**

2    **Participants**

3            We included 4,529 participants without any missing data pertaining to soda intake, risk  
4    factors for SUDs, alcohol sipping, and all the confounding variables. The participants were  
5    divided into four groups based on their caffeinated soda consumption: the non-drinking group  
6    ( $N=1,945$ ), monthly drinking group ( $N=1,941$ ), weekly drinking group ( $N=484$ ), and daily drinking  
7    group ( $N=159$ ) [36]. See **Supplementary Materials** for more details on the measures and  
8    participants.

9

10    **Analysis**

11    *Caffeinated soda intake and risk factors for SUDs*

12           To identify a multivariate profile of risk factors for caffeinated soda consumption, we  
13    performed a least absolute shrinkage and selection operator (LASSO) regression analysis [37], a  
14    machine learning algorithm, to distinguish the daily drinking group ( $N=159$ ) from the non-drinking  
15    group ( $N=1,945$ ). We focused on these two extreme groups to ensure adequate separation  
16    between habitual drinkers (daily drinking group) and non-drinkers (non-drinking group). We used  
17    all of the risk factors and confounding variables as input (i.e., predictors) in the LASSO-based  
18    prediction model to classify each individual into the daily drinking group (coded as "1") or non-  
19    drinking group (coded as "0"). More specifically, candidate predictors included every measure for  
20    the three cognitive factors as well as other control variables. See **Supplementary Materials** for  
21    more details on the predictors and LASSO regression.

22

23    *Caffeinated soda intake and future alcohol sipping*

24           To rigorously examine the association between caffeinated soda intake and future  
25    alcohol sipping, we used univariate (hierarchical regression) and multivariate (machine learning)  
26    approaches. Hierarchical logistic regression analysis was used to test whether caffeinated soda  
27    intake at baseline can predict alcohol sipping after 12 months. To ensure that the main predictor

1 accounted for more variance than the confounding variables, we set a three-level hierarchical  
2 model (**Table 1**). See **Supplementary Materials** for more details on the hierarchical regression.

3 We also performed multivariate LASSO regression analysis to confirm the association  
4 between caffeinated soda intake and alcohol sipping. We used all possible variables, including  
5 the three neurocognitive risk factors (WM, impulsivity, and reward processing) and the  
6 confounding variables as candidate predictors collected at baseline. Using these inputs (i.e.,  
7 predictors) and the LASSO-based model, we aimed to classify individuals who reported alcohol  
8 sipping after 12 months (coded as "1") and those who reported no alcohol sipping (coded as "0").  
9 All of the procedures, except for the predictors and dependent variables, are identical to those in  
10 the former analysis (see **METHODS - Caffeinated soda intake and risk factors for SUDs**).

11

12 *Caffeinated soda intake, alcohol sipping, and risk factors for SUDs*

13 We performed the mediation analysis to provide a more integrated insight into how the  
14 interactions between caffeinated soda intake and the SUD risk factors lead to future alcohol  
15 sipping in children. We fit three mediation models for each SUD risk factor, with a mediation  
16 design involving the following four elements: an independent variable (one neurocognitive risk  
17 factor for one model - WM, impulsivity, and reward processing), two mediators (caffeinated soda  
18 intake and family risk factors; covariance of these two mediators was included in the model), and  
19 a dependent variable (alcohol sipping after 12 months). This design was chosen to identify a  
20 neurocognitive risk factor that is mediated by caffeinated soda intake in predicting alcohol sipping.

21 See **Supplementary Materials** for more details on the mediation analysis.

22

## 23 **RESULTS**

24 **Caffeinated soda intake and risk factors for SUDs**

25 Our first question was whether caffeinated soda intake is associated with well-known  
26 neurocognitive risk factors for SUDs (**Figure 1A**). **Figure 2A** shows the multivariate profiles from  
27 the LASSO regression distinguishing the daily-drinking group from the non-drinking group. Sex  
28 (being male) and low parental education were the two strongest predictors of daily consumption,

1 along with a family history of drug use, high Behavioral Activation System (BAS) scores [38], lack  
2 of sleep, and low family income. **Figure 2B** shows a receiver-operation characteristic (ROC)  
3 curve and its mean area under the curve (AUC) for classification of the daily drinking and non-  
4 drinking groups. The mean AUC values were 0.82 and 0.73 for the training and test sets,  
5 respectively.

6 Among the risk factors for SUDs, high impulsivity measured using the BAS was the most  
7 predictive of daily caffeinated soda intake (**Figure 2C**). A higher Urgency–Premeditation–  
8 Perseverance–Sensation Seeking–Positive Urgency (UPPS-P) impulsive behavior score [39] and  
9 a lower Behavioral Inhibition System (BIS) score [38] were related to daily intake. Additionally,  
10 hyperactivation in the nucleus accumbens (NAc) and pars triangularis in the inferior frontal gyrus  
11 (IFG) and hypoactivation in the caudal anterior cingulate cortex (ACC) and caudate nucleus  
12 during the stop signal task (SST) predicted the classification of the daily drinking group. Among  
13 the WM measures, poor performance on the list sorting test, card sort test, and 0-back  
14 performance in the emotional N-Back (EN-back) task predicted daily intake of caffeinated soda. In  
15 addition, hypoactivation of the pars orbitalis of the IFG and greater activation of the NAc, caudate  
16 nucleus, inferior parietal lobule (IPL), and thalamus proper during the EN-back task (2-back vs. 0-  
17 back) also predicted daily caffeinated soda intake (**Figure 2C**). The daily drinking group showed a  
18 smaller difference in accuracy between the reward and neutral trials on the monetary incentive  
19 delay (MID) task compared to the non-drinking group, suggesting reduced motivation to achieve  
20 the monetary reward. Additionally, hyperactivity in the medial orbitofrontal cortex (OFC) and  
21 hypoactivity in the rostral ACC during the MID task (reward vs. neutral anticipation) predicted the  
22 classification into the daily drinking group.

23 These results support that low WM, high impulsivity, and blunted reward anticipation are  
24 significantly associated with daily caffeinated soda consumption. Along with some other  
25 demographic factors (male sex, family history of drug use, low socioeconomic status (SES), low  
26 parental monitoring, high externalizing behaviors, less sleep, high body mass index (BMI), older  
27 age, and low physical activity), the three neurocognitive risk factors for SUDs distinguished the  
28 daily drinking group from the non-drinking group.

1     *Caffeinated soda intake and future alcohol sipping*

2           We first conducted a three-level hierarchical logistic regression analysis to address our  
3     second main question of whether caffeinated soda intake predicts future alcohol sipping, (**Table**  
4     **1**). Specifically, we examined whether caffeinated soda intake accounted for additional variance  
5     than did other confounding variables. The adjusted R-square value of Model 3 revealed a  
6     significant difference compared with Model 1 ( $F(4, 8,854)=177.58$ ,  $p<2.20e-16$ ) and Model 2  
7     ( $F(3,8,854)=6.61$ ,  $p<0.001$ ), supporting the hypothesis that caffeinated soda drinking behavior in  
8     childhood can predict future alcohol drinking (**Figure 3A**). Higher caffeinated soda consumption  
9     significantly predicted alcohol sipping after 12 months even after controlling for confounding  
10   variables and baseline alcohol sipping (**Table 1**).

11          Next, we performed LASSO regression predicting alcohol sipping after 12 months, using  
12     all of the neurocognitive risk factors and confounding variables collected at baseline. As shown in  
13     **Figure 3B**, caffeinated soda intake (mean estimate of coefficients=0.138, 95% confidence  
14     interval (CI)=[0.123, 0.155]) was identified as a predictor that distinguished alcohol sipping after  
15     12 months, even after including alcohol sipping at baseline as predictors. The mean AUC values  
16     of the predictive model were 0.81 and 0.76 for the training and test sets, respectively (**Figure 3C**).  
17

18     *Caffeinated soda intake, risk factors for SUDs, and future alcohol sipping*

19          We examined the integrated relationships among the risk factors for SUDs, caffeinated  
20     soda intake, and alcohol sipping after 12 months. We tested three mediation models using each  
21     neurocognitive risk factor (WM, impulsivity, and reward processing) as the independent variable.  
22     The models that used WM or reward processing as an independent variable showed inconclusive  
23     total effects (WM model:  $B=-0.007$ , 95% CI=[−0.013. −0.001],  $p=0.232$ ; reward processing  
24     model:  $B=0.004$ , 95% CI=[−0.002. 0.010],  $p=0.457$ ). However, as shown in **Figure 4**, the model  
25     using impulsivity as an independent variable revealed significant total effects ( $B=0.023$ , 95%  
26     CI=[0.018, 0.028],  $p<0.001$ ). The model detected significant indirect effects of impulsivity on  
27     alcohol sipping after 12 months, with caffeinated soda intake and family risk factors mediating the  
28     relationships (caffeinated soda intake:  $B=0.002$ , 95% CI=[0.001, 0.003],  $p=0.013$ ; family risk

1 factor:  $B=-0.002$ , 95% CI=[-0.003, -0.001],  $p=0.006$ ). These results suggest caffeinated soda  
2 intake mediates impulsivity as the SUD risk factors toward future alcohol sipping.

3

#### 4 **DISCUSSION**

5 In this study, we investigated whether caffeinated soda intake in childhood is associated  
6 with a higher risk of alcohol use in the future. Using the large dataset from the ABCD study  
7 ( $N=4,529$ ), we identified the relationships between caffeinated soda intake and the neurocognitive  
8 risk factors for SUDs, evaluated the predictive ability of caffeinated soda on alcohol sipping after  
9 12 months, and tested the hypothesis that caffeinated soda intake mediates neurocognitive risk  
10 factors and alcohol sipping after 12 months. Our findings suggest that frequent consumption of  
11 caffeinated soda in children is closely related to the neurocognitive risk factors for SUDs and can  
12 predict future alcohol sipping, particularly by mediating impulsivity.

13 The machine learning approach revealed a strong association between caffeinated soda  
14 intake and the neurocognitive risk factors for SUDs. High impulsivity scores, low WM  
15 performance, and blunted responses to monetary reward distinguished daily caffeinated soda  
16 drinkers from non-drinkers (**Figure 2**). In particular, we showed higher self-reported impulsivity in  
17 the daily drinking group based on the UPPS-P and BIS/BAS scores along with altered activation  
18 in regions implicated in impulsivity, i.e., hypoactivity in the ACC and caudate nucleus and  
19 hyperactivity in the IFG and NAc during the SST. Consistent with our findings, reduced activities  
20 in the ACC and caudate nucleus are commonly reported in children with attention deficit  
21 hyperactivity disorder (ADHD) during response inhibition tasks [40–44], and activation of the  
22 ventral striatum is greater in individuals with SUDs, implying dysregulation of impulsivity [45–47].  
23 We detected hyperactivity in the IFG among daily soda drinkers compared with non-drinkers.  
24 Previous studies have reported reduced activity in the IFG as neural markers of substance abuse  
25 and ADHD [48–50]. However, increased activity in the IFG has also been reported in individuals  
26 with tobacco use disorder [51], and increased IFG activity suggests that more preparatory effort is  
27 required for successful inhibition [52].

1 We also showed behavioral WM impairments in the daily drinking group on the list sorting  
2 test, card sort test, and the 0-back accuracy in the EN-back task, accompanied by hypoactivation  
3 in the IFG and hyperactivation in the NAc, caudate nucleus, IPL, and thalamus during the EN-  
4 back task. These results are consistent with the developmental mismatch hypothesis, which  
5 states that subcortical structures of affect and reward processing develop earlier than prefrontal  
6 structures involved in the top-down regulation of behavior in children and adolescents [53,54].  
7 Based on the previous studies that cognitive control during the EN-back task is related to WM  
8 capacity in the prefrontal cortex and simultaneous emotional regulation in the limbic system  
9 [30,55], the low WM performance in the daily drinking group may be related to increased  
10 activation in the limbic system and decreased activation in the IFG activation during the WM task.

11 Blunted reward anticipation was associated with caffeinated soda intake. Along with a  
12 smaller motivation to achieve the reward, activation in the ACC was lower and activation in the  
13 medial OFC was higher during the MID task in the daily drinking group than in the non-drinking  
14 group. Hyperactivation in the medial prefrontal cortex (mPFC) regions during reward anticipation  
15 is supported by previous literature suggesting increased prefrontal region activity, although these  
16 findings do not always replicate [56–60].

17 We showed that frequent consumption of caffeinated soda predicted alcohol sipping after  
18 12 months using hierarchical regression (**Table 1**) and LASSO regression (**Figure 3**). Among the  
19 significant predictors in the LASSO regression, caffeinated soda intake was the second strongest  
20 predictor of future alcohol sipping, following the baseline alcohol sipping (**Figure 3B**). Other  
21 survived predictors include predictors for daily caffeinated soda intake (**Figure 2A**) while others  
22 showed the opposite direction in predicting daily caffeinated soda intake and alcohol sipping after  
23 12 months. For example, higher SES, greater physical activity, no family history of drug use,  
24 younger age, and reduced activity in the ACC during the EN-back task and in the medial OFC  
25 during the MID task were associated with the participants who experienced alcohol sipping.  
26 Interestingly, predictors of caffeinated soda intake were similar to the risk factors for SUDs  
27 [18,61], suggesting that caffeinated soda intake during childhood and SUDs share similar  
28 neurocognitive vulnerabilities. Note that the majority of the cohort was substance-naïve [62], and

1 we evaluated alcohol sipping measures instead of actual administration of the substance. Thus, it  
2 would be informative to track the predictive ability of caffeinated soda and alcohol use and  
3 investigate changes in the direction of the predictors as the children get older.

4 The mediation analysis revealed that caffeinated soda consumption is a mediator  
5 between impulsivity and future alcohol sipping (**Figure 4**). This finding suggests that caffeinated  
6 soda itself may not directly cause children to initiate substance use, but the drinking behavior of  
7 the beverages could indicate high impulsivity, which may be linked to a high risk of initiating  
8 substance use in the near future. Because caffeinated soda contains two addictive substances,  
9 sugar and caffeine, it is somewhat natural for children to prefer and seek the beverages. While  
10 this taste preference could lead to a seeking behavior and habitual consumption, previous studies  
11 have shown impulsivity as one of the most common traits of SUDs [63–65]. Therefore, the  
12 behavior of frequently consuming caffeinated soda could indicate a high risk of initiating  
13 substance use in the future, although further research is needed to disentangle this complex  
14 relationship. Along with caffeinated soda intake, family risk factors also mediated the relationship  
15 between impulsivity and future alcohol sipping; higher impulsivity was associated with having  
16 more family risk factors, which led to less alcohol sipping after 12 months. Notably, some  
17 previous studies suggested that family risk factors, such as low SES, increase the risk of future  
18 alcohol use [66–69], while others suggested the opposite [70,71].

19 To our knowledge, this is the first study to investigate the direct link between caffeinated  
20 soda intake in childhood and the risk of substance use. The results are consistent with studies on  
21 caffeine consumption during adolescence and its association with future substance misuse  
22 [10,13], supporting a higher risk of caffeinated soda consumption in childhood, particularly  
23 regarding vulnerability to future substance misuse. Such information is invaluable, as caffeinated  
24 soda is incomparably the most common medium for caffeine consumption in childhood, and the  
25 risk of substance misuse should be detected before adolescence, the most common period of  
26 substance use onset.

27 Our results have important implications for public health recommendations, as our study  
28 provides novel insight into the neurocognitive correlates of caffeinated soda consumption in

1 children, which has rarely been evaluated. At the same time, a few limitations of our research and  
2 future directions for further investigations should be discussed. First, we did not perform  
3 functional connectivity analyses or multivoxel pattern analyses, which may have provided  
4 additional insight into the effects of caffeinated soda intake. Second, as the 9–10-year-old  
5 children in this study had not yet started other substances except for alcohol, such as tobacco or  
6 marijuana, future work using the longitudinal 10-year follow-up data of the ABCD study should  
7 examine whether frequent consumption of caffeinated soda is associated with alcohol or other  
8 substance misuse. Third, we acknowledge that multiple variables other than caffeinated soda  
9 intake and family risk factors may mediate the relationship between neurocognitive risk factors  
10 and future alcohol use; thus, extensive investigation of how caffeinated soda intake interacts with  
11 other SUD risk/protective factors is needed in the future. Lastly, the ABCD dataset included only  
12 a small set of measures of food/drink consumption. Future studies would benefit by including  
13 more diverse measures of foods or drinks containing caffeine or sugar; for example, by  
14 determining if similar results are found with non-caffeinated soda or other sugary beverages.

15 In conclusion, our results revealed the potential risks of caffeinated soda consumption in  
16 children by investigating the associations between caffeinated soda consumption and risk factors  
17 for SUDs, examining the ability of caffeinated soda consumption to predict future alcohol sipping,  
18 and revealing that caffeinated soda consumption mediates impulsivity and future alcohol sipping,  
19 using the large ABCD dataset. While previous research on the side effects of caffeinated soda  
20 consumption has been limited to negative physical consequences, the present results strongly  
21 suggest that caffeinated soda drinking in children is also associated with changes in  
22 neurocognitive function and can predict alcohol sipping after 12 months. Our study further  
23 suggests a strong need to develop evidence-based recommendations for caffeinated soda [2], as  
24 there is no consensus on a safe dose of caffeinated soda in children, and some children are at  
25 higher risk of adverse events from frequent caffeinated soda intake. Further clarification on the  
26 causal relationships and neuro-developmental evidence are needed to determine whether  
27 caffeinated soda is a warning sign for future substance misuse and whether it induces  
28 neurocognitive impairments in children.

## 1   **ACKNOWLEDGEMENTS**

2   We thank the families and children who participated, and continue to participate, in the ABCD  
3   study, as well as staff at the study sites who are involved in data collection and curating the data  
4   release. Data used in the preparation of this article were obtained from the Adolescent Brain  
5   Cognitive Development (ABCD) Study (<https://abcdstudy.org>), held in the NIMH Data Archive  
6   (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-  
7   10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the  
8   National Institutes of Health and additional federal partners under award numbers U01DA041048,  
9   U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987,  
10   U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988,  
11   U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148,  
12   U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is  
13   available at <https://abcdstudy.org/federal-partners.html>. A listing of participating sites and a  
14   complete listing of the study investigators can be found at  
15   [https://abcdstudy.org/consortium\\_members/](https://abcdstudy.org/consortium_members/). ABCD consortium investigators designed and  
16   implemented the study and/or provided data but did not necessarily participate in analysis or  
17   writing of this report. This manuscript reflects the views of the authors and may not reflect the  
18   opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows  
19   and changes over time. The ABCD data used in this report came from [NIMH Data Archive Digital  
20   Object Identifier (10.15154/1503209)]. DOIs can be found at <http://dx.doi.org/10.15154/1503209>.

## REFERENCES

- 1
- 2 1. Temple JL. Caffeine use in children: What we know, what we have left to learn, and why we  
3 should worry. *Neurosci Biobehav Rev.* 2009;33(6):793–806.
- 4 2. Temple JL. Trends, Safety, and Recommendations For Caffeine Use in Children and  
5 Adolescents. *J Am Acad Child Adolesc Psychiatry.* 2018;58(1):36–45.
- 6 3. Ooi JY, Wolfenden L, Sutherland R, Nathan N, Oldmeadow C, McLaughlin M, et al. A  
7 Systematic Review of the Recent Consumption Levels of Sugar-Sweetened Beverages in  
8 Children and Adolescents From the World Health Organization Regions With High Dietary–  
9 Related Burden of Disease. *Asia Pac J Public Heal.* 2022;34(1):11–24.
- 10 4. Sylvetsky AC, Parnarouskis L, Merkel PE, Gearhardt AN. Children’s Sugar-Sweetened  
11 Beverage Consumption: Striking Parallels With Substance Use Disorder Symptoms. *Frontiers  
12 Pediatrics.* 2020;8:594513.
- 13 5. Nehlig A. Are we dependent upon coffee and caffeine? A review on human and animal data.  
14 *Neurosci Biobehav Rev.* 1999;23(4):563–76.
- 15 6. Avena NM, Rada P, Hoebel BG. Evidence for sugar addiction: Behavioral and neurochemical  
16 effects of intermittent, excessive sugar intake. *Neurosci Biobehav Rev.* 2008;32(1):20–39.
- 17 7. Porciúncula LO, Sallaberry C, Mioranza S, Botton PHS, Rosemberg DB. The Janus face of  
18 caffeine. *Neurochem Int.* 2013;63(6):594–609.
- 19 8. Arria AM, Caldeira KM, Kasperski SJ, Vincent KB, Griffiths RR, O’Grady KE. Energy Drink  
20 Consumption and Increased Risk for Alcohol Dependence. *Alcohol Clin Exp Res.*  
21 2011;35(2):365–75.
- 22 9. Miyake ER, Marmorstein NR. Energy drink consumption and later alcohol use among early  
23 adolescents. *Addict Behav.* 2015;43:60–5.
- 24 10. Barrense-Dias Y, Berchtold A, Akre C, Suris J. Consuming energy drinks at the age of 14  
25 predicted legal and illegal substance use at 16. *Acta Paediatr.* 2016;105(11):1361–8.
- 26 11. Kristjansson AL, Kogan SM, Mann MJ, Smith ML, Juliano LM, Lilly CL, et al. Does early  
27 exposure to caffeine promote smoking and alcohol use behavior? A prospective analysis of  
28 middle school students. *Addiction.* 2018;113(9):1706–13.
- 29 12. Marmorstein NR. Investigating associations between caffeinated beverage consumption and  
30 later alcohol consumption among early adolescents. *Addict Behav.*  
31 2018;90(Psychopharmacology 203 2009):362–8.
- 32 13. Leal WE, Jackson DB. The role of energy drink consumption in the intention to initiate  
33 marijuana use among adolescents. *Addict Behav.* 2019;93(Postgraduate Medicine 127 3  
34 2015):240–5.

- 1    14. Lisdahl KM, Tapert S, Sher KJ, Gonzalez R, Nixon SJ, Ewing SWF, et al. Substance use  
2    patterns in 9-10 year olds: Baseline findings from the adolescent brain cognitive development  
3    (ABCD) study. *Drug Alcohol Depen.* 2021;108946.
- 4    15. Vercammen KA, Koma JW, Bleich SN. Trends in Energy Drink Consumption Among U.S.  
5    Adolescents and Adults, 2003–2016. *Am J Prev Med.* 2019;56(6):827–33.
- 6    16. Chambers RA, Taylor JR, Potenza MN. Developmental Neurocircuitry of Motivation in  
7    Adolescence: A Critical Period of Addiction Vulnerability. *Am J Psychiat.* 2003;160(6):1041–52.
- 8    17. Wagner FA, Anthony JC. From First Drug Use to Drug Dependence: Developmental Periods  
9    of Risk for Dependence upon Marijuana, Cocaine, and Alcohol. *Neuropsychopharmacol.*  
10   2002;26(4):479–88.
- 11   18. Jordan CJ, Andersen SL. Sensitive periods of substance abuse: Early risk for the transition to  
12   dependence. *Dev Cogn Neurosci.* 2017;25:29–44.
- 13   19. Barry AE, King J, Sears C, Harville C, Bondoc I, Joseph K. Prioritizing Alcohol Prevention:  
14   Establishing Alcohol as the Gateway Drug and Linking Age of First Drink With Illicit Drug Use. *J  
15   School Health.* 2016;86(1):31–8.
- 16   20. Schneider S, Schilling L, Osenbrügge N. Determinants of soft drink consumption among  
17   children and adolescents in developed countries - a systematic review. *Cent Eur J Publ Heal.*  
18   2021;29(4):290–300.
- 19   21. Lopez NV, Ayala GX, Corder K, Eisenberg CM, Zive MM, Wood C, et al. Parent Support and  
20   Parent-Mediated Behaviors Are Associated with Children’s Sugary Beverage Consumption. *J  
21   Acad Nutr Diet.* 2012;112(4):541–7.
- 22   22. Zahid A, Davey C, Reicks M. Beverage Intake among Children: Associations with Parent and  
23   Home-Related Factors. *Int J Environ Res Pu.* 2017;14(8):929.
- 24   23. Gilligan C, Kyri K. Parent attitudes, family dynamics and adolescent drinking: qualitative  
25   study of the Australian parenting guidelines for adolescent alcohol use. *Bmc Public Health.*  
26   2012;12(1):491–491.
- 27   24. Underwood JM, Brener N, Thornton J, Harris WA, Bryan LN, Shanklin S, et al. Youth Risk  
28   Behavior Surveillance — United States, 2019. Centers for Disease Control and Prevention. 2020  
29   Aug 21;
- 30   25. Muchiri BW, Santos MML dos. Family management risk and protective factors for adolescent  
31   substance use in South Africa. *Subst Abus Treat Prev Policy.* 2018;13(1):24.
- 32   26. Wills TA, Yaeger AM. Family Factors and Adolescent Substance Use. *Curr Dir Psychol Sci.*  
33   2003;12(6):222–6.
- 34   27. Solnick SJ, Hemenway D. Soft drinks, aggression and suicidal behaviour in US high school  
35   students. *Int J Inj Control Sa.* 2013;21(3):266–73.
- 36   28. James JE, Kristjánsson ÁL, Sigríðsdóttir ID. Adolescent substance use, sleep, and academic  
37   achievement: Evidence of harm due to caffeine. *J Adolescence.* 2011;34(4):665–73.

- 1    29. Suglia SF, Solnick S, Hemenway D. Soft Drinks Consumption Is Associated with Behavior  
2    Problems in 5-Year-Olds. *J Pediatrics*. 2013;163(5):1323–8.
- 3    30. Casey BJ, Cannonier T, Conley MI, Cohen AO, Barch DM, Heitzeg MM, et al. The Adolescent  
4    Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn  
5    Neurosci*. 2018;32(Biol. Psychiatry 69 2011):43–54.
- 6    31. Volkow ND, Koob G, Baler R. Biomarkers in Substance Use Disorders. *Acs Chem Neurosci*.  
7    2015;6(4):522–5.
- 8    32. Watts AL, Wood PK, Jackson KM, Lisdahl KM, Heitzeg MM, Gonzalez R, et al. Incipient  
9    alcohol use in childhood: Early alcohol sipping and its relations with psychopathology and  
10   personality. *Dev Psychopathol*. 2021;33(4):1338–50.
- 11   33. Jackson KM, Barnett NP, Colby SM, Rogers ML. The Prospective Association Between  
12   Sipping Alcohol by the Sixth Grade and Later Substance Use. *J Stud Alcohol Drugs*.  
13   2015;76(2):212–21.
- 14   34. Donovan JE, Molina BSG. Childhood Risk Factors for Early-Onset Drinking\*. *J Stud Alcohol  
15   Drugs*. 2011;72(5):741–51.
- 16   35. Kirby T, Barry AE. Alcohol as a gateway drug: a study of US 12th graders. *J Sch Heal*.  
17   2012;82(8):371–9.
- 18   36. Mullee A, Romaguera D, Pearson-Stuttard J, Viallon V, Stepien M, Freisling H, et al.  
19   Association Between Soft Drink Consumption and Mortality in 10 European Countries. *Jama  
20   Intern Med*. 2019;179(11).
- 21   37. Tibshirani R. Regression Shrinkage and Selection Via the Lasso. *J Royal Statistical Soc Ser  
22   B Methodol*. 1996;58(1):267–88.
- 23   38. Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to  
24   impending reward and punishment: The BIS/BAS Scales. *J Pers Soc Psychol*. 1994;67(2):319.
- 25   39. Barch DM, Belden AC, Tillman R, Whalen D, Luby JL. Early Childhood Adverse Experiences,  
26   Inferior Frontal Gyrus Connectivity, and the Trajectory of Externalizing Psychopathology. *J Am  
27   Acad Child Adolesc Psychiatry*. 2018;57(3):183–90.
- 28   40. Lei D, Du M, Wu M, Chen T, Huang X, Du X, et al. Functional MRI Reveals Different  
29   Response Inhibition Between Adults and Children With ADHD. *Neuropsychology*.  
30   2015;29(6):874–81.
- 31   41. Cortese S, Kelly C, Chabernaud C, Proal E, Martino AD, Milham MP, et al. Toward Systems  
32   Neuroscience of ADHD: A Meta-Analysis of 55 fMRI Studies. *Am J Psychiatr*. 2012;169(10):1038–  
33   55.
- 34   42. Hart H, Radua J, Nakao T, Mataix-Cols D, Rubia K. Meta-analysis of Functional Magnetic  
35   Resonance Imaging Studies of Inhibition and Attention in Attention-deficit/Hyperactivity Disorder:  
36   Exploring Task-Specific, Stimulant Medication, and Age Effects. *Jama Psychiatr*. 2013;70(2):185–  
37   98.

- 1    43. Zarka D, Cebolla AM, Cevallos C, Palmero-Soler E, Dan B, Cheron G. Caudate and  
2    cerebellar involvement in altered P2 and P3 components of GO/NoGO evoked potentials in  
3    children with attention-deficit/hyperactivity disorder. *Eur J Neurosci*. 2021;53(10):3447–62.
- 4    44. Mahmood OM, Goldenberg D, Thayer R, Migliorini R, Simmons AN, Tapert SF. Adolescents'  
5    fMRI activation to a response inhibition task predicts future substance use. *Addict Behav*.  
6    2013;38(1):1435–41.
- 7    45. Bjork JM, Smith AR, Hommer DW. Striatal sensitivity to reward deliveries and omissions in  
8    substance dependent patients. *Neuroimage*. 2008;42(4):1609–21.
- 9    46. Volkow ND, Fowler JS, Wang GJ, Swanson JM, Telang F. Dopamine in Drug Abuse and  
10   Addiction: Results of Imaging Studies and Treatment Implications. *Arch Neurol-chicago*.  
11   2007;64(11):1575–9.
- 12   47. Heitzeg MM, Nigg JT, Yau WYW, Zucker RA, Zubieta JK. Striatal Dysfunction Marks  
13   Preexisting Risk and Medial Prefrontal Dysfunction Is Related to Problem Drinking in Children of  
14   Alcoholics. *Biol Psychiat*. 2010;68(3):287–95.
- 15   48. Schmidt A, Walter M, Gerber H, Schmid O, Smieskova R, Bendfeldt K, et al. Inferior Frontal  
16   Cortex Modulation with an Acute Dose of Heroin During Cognitive Control.  
17   *Neuropsychopharmacol*. 2013;38(11):2231–9.
- 18   49. Fu L ping, Bi G hua, Zou Z tong, Wang Y, Ye E mao, Ma L, et al. Impaired response inhibition  
19   function in abstinent heroin dependents: An fMRI study. *Neurosci Lett*. 2008;438(3):322–6.
- 20   50. Morein-Zamir S, Dodds C, Harteveld TJ, Schwarzkopf W, Sahakian B, Müller U, et al.  
21   Hypoactivation in right inferior frontal cortex is specifically associated with motor response  
22   inhibition in adult ADHD. *Hum Brain Mapp*. 2014;35(10):5141–52.
- 23   51. Kalhan S, Chen LPE, Garrido MI, Hester R. People with tobacco use disorder exhibit more  
24   prefrontal activity during preparatory control but reduced anterior cingulate activity during reactive  
25   control. *Addict Biol*. 2022;27(2):e13159.
- 26   52. DeVito EE, Meda SA, Jantionio R, Potenza MN, Krystal JH, Pearlson GD. Neural Correlates  
27   of Impulsivity in Healthy Males and Females with Family Histories of Alcoholism.  
28   *Neuropsychopharmacol*. 2013;38(10):1854–63.
- 29   53. Somerville LH, Hare T, Casey BJ. Frontostriatal maturation predicts cognitive control failure to  
30   appetitive cues in adolescents. *J Cognitive Neurosci*. 2010;23(9):2123–34.
- 31   54. Mills KL, Goddings AL, Clasen LS, Giedd JN, Blakemore SJ. The Developmental Mismatch in  
32   Structural Brain Maturation during Adolescence. *Dev Neurosci-basel*. 2014;36(3–4):147–60.
- 33   55. Gee DG, Humphreys KL, Flannery J, Goff B, Telzer EH, Shapiro M, et al. A developmental  
34   shift from positive to negative connectivity in human amygdala-prefrontal circuitry. *J Neurosci*  
35   Official J Soc Neurosci. 2013;33(10):4584–93.
- 36   56. Beck A, Schlagenhauf F, Wüstenberg T, Hein J, Kienast T, Kahnt T, et al. Ventral Striatal  
37   Activation During Reward Anticipation Correlates with Impulsivity in Alcoholics. *Biol Psychiat*.  
38   2009;66(8):734–42.

- 1        57. Balodis IM, Kober H, Worhunsky PD, Stevens MC, Pearlson GD, Potenza MN. Diminished  
2        Frontostriatal Activity During Processing of Monetary Rewards and Losses in Pathological  
3        Gambling. *Biol Psychiat*. 2012;71(8):749–57.
- 4        58. Swartz JR, Weissman DG, Ferrer E, Beard SJ, Fassbender C, Robins RW, et al. Reward-  
5        Related Brain Activity Prospectively Predicts Increases in Alcohol Use in Adolescents. *J Am Acad*  
6        *Child Adolesc Psychiatry*. 2019;59(3):391–400.
- 7        59. Elsayed NM, Kim MJ, Fields KM, Olvera RL, Hariri AR, Williamson DE. Trajectories of Alcohol  
8        Initiation and Use During Adolescence: The Role of Stress and Amygdala Reactivity. *J Am Acad*  
9        *Child Adolesc Psychiatry*. 2018;57(Environ Psychol Nonverbal Behav 1 1 1976):550–60.
- 10      60. Luijten M, Schellekens AF, Kühn S, Machielse MWJ, Sescousse G. Disruption of Reward  
11      Processing in Addiction : An Image-Based Meta-analysis of Functional Magnetic Resonance  
12      Imaging Studies. *Jama Psychiat*. 2017;74(4):387.
- 13      61. Kilpatrick DG, Acierno R, Saunders B, Resnick HS, Best CL, Schnurr PP. Risk Factors for  
14      Adolescent Substance Abuse and Dependence: Data From a National Sample. *J Consult Clin*  
15      *Psych*. 2000;68(1):19–30.
- 16      62. Lisdahl KM, Sher KJ, Conway KP, Gonzalez R, Ewing SWF, Nixon SJ, et al. Adolescent brain  
17      cognitive development (ABCD) study: Overview of substance use assessment methods. *Dev*  
18      *Cogn Neurosci*. 2018;32:80–96.
- 19      63. Cyders MA, Flory K, Rainer S, Smith GT. The role of personality dispositions to risky behavior  
20      in predicting first-year college drinking. *Addiction*. 2009;104(2):193–202.
- 21      64. Whiteside SP, Lynam DR. Understanding the Role of Impulsivity and Externalizing  
22      Psychopathology in Alcohol Abuse: Application of the UPPS Impulsive Behavior Scale. *Exp Clin*  
23      *Psychopharm*. 2003;11(3):210–7.
- 24      65. Magid V, Colder CR. The UPPS Impulsive Behavior Scale: Factor structure and associations  
25      with college drinking. *Pers Indiv Differ*. 2007;43(7):1927–37.
- 26      66. Calling S, Ohlsson H, Sundquist J, Sundquist K, Kendler KS. Socioeconomic status and  
27      alcohol use disorders across the lifespan: A co-relative control study. *Plos One*.  
28      2019;14(10):e0224127.
- 29      67. Carroll HA, Heleniak C, Witkiewitz K, Lewis M, Eakins D, Staples J, et al. Effects of parental  
30      monitoring on alcohol use in the US and Sweden: A brief report. *Addict Behav*. 2016;63:89–92.
- 31      68. Collins SE. Associations Between Socioeconomic Factors and Alcohol Outcomes. *Alcohol Res Curr Rev*. 2016;38(1):83–94.
- 33      69. Goodwin DW. Alcoholism and Heredity: A Review and Hypothesis. *Arch Gen Psychiat*.  
34      1979;36(1):57–61.
- 35      70. Liu Y, Lintonen T, Tynjälä J, Villberg J, Välimäa R, Ojala K, et al. Socioeconomic differences in  
36      the use of alcohol and drunkenness in adolescents: Trends in the Health Behaviour in School-  
37      aged Children study in Finland 1990–2014. *Scand J Public Healt*. 2016;46(1):102–11.


1    71. Kendler KS, Gardner CO, Hickman M, Heron J, Macleod J, Lewis G, et al. Socioeconomic  
2    Status and Alcohol-Related Behaviors in Mid- to Late Adolescence in the Avon Longitudinal  
3    Study of Parents and Children. *J Stud Alcohol Drugs*. 2014;75(4):541–5.

4    72. Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri and  
5    sulci using standard anatomical nomenclature. *Neuroimage*. 2010;53(1):1–15.

6

1    **Figure 1.** An overview of the study design

2    **A.** Diagram of the research aims. First, the association between the risk factors for substance use  
3    disorders (SUDs) and caffeinated soda intake was examined. Then, we assessed whether  
4    caffeinated soda intake predicts alcohol sipping after 12 months, and whether caffeinated soda  
5    intake works as a mediator between the risk factors and future alcohol sipping. **B.** The variables  
6    used to measure the three main neurocognitive risk factors: working memory, impulsivity, and  
7    reward processing. Working memory was measured from the performance of list sorting test, card  
8    sort test, and 2-back in the emotional N-Back Task (EN-back) performed inside a functional  
9    magnetic resonance imaging (fMRI) scanner. The fMRI data from the EN-back task were  
10   analyzed based on the contrast of 2-back versus 0-back, using regions in the frontoparietal and  
11   fronto-thalamic network as the regions of interest (ROIs). Impulsivity was measured by Urgency-  
12   Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P), and Behavioral  
13   Inhibition System/ Behavioral Activation System (BIS/BAS), and the stop signal reaction time  
14   (SSRT) during the stop signal task (SST) performed inside the fMRI scanner. The fMRI data from  
15   the SST were analyzed based on the contrast of correct stop versus correct go, using regions in  
16   the lateral prefrontal cortex, anterior cingulate cortex (ACC), and striatum as the ROIs. Reward  
17   processing was measured by cash choice task, and also by comparing the success rate of  
18   reward versus neutral conditions during the monetary incentive delay (MID) task performed inside  
19   the fMRI scanner. The fMRI data from the MID task were analyzed based on the contrast of  
20   reward anticipation versus neutral anticipation, using regions in the ventral striatum and medial  
21   frontal cortex as the ROIs. The ROIs were selected based on the Destrieux atlas [72].




1

1   **Figure 2.** Results of LASSO regression predicting daily caffeinated soda intake (daily drinking  
2   group vs. non-drinking group)

3   **A.** Estimates of coefficients of the survived variables (x axis: coefficient estimates; y axis: predictor).  
4   The site variables were excluded for clarify (see **Figure S1** for the site variables). **B.** Distribution of  
5   the area under the curve (AUC) values (left) and a representative receiver-operation characteristic  
6   (ROC) curve (right) for the training and test datasets. **C.** Regions of interest (ROIs) identified as  
7   having significant estimates of coefficients during the emotional N-Back Task (EN-back), stop signal  
8   task (SST), and monetary incentive delay (MID) task.


9   *Abbreviations.* LASSO, least absolute shrinkage and selection operator; BMI, body mass index;  
10   BIS, Behavioral Inhibition System; BAS, Behavioral Activation System; UPPS-P, Urgency-  
11   Premarkption-Perseverance-Sensation Seeking-Positive Urgency; SST, Stop Signal Task; SSRT,  
12   Stop Signal Reaction Time; MID, Monetary Incentive Delay Task; NAc, Nucleus Accumbens; IPL,  
13   Inferior Parietal Lobule; IFG, Inferior Frontal Gyrus; ACC, Anterior Cingulate Cortex; OFC,  
14   Orbitofrontal Cortex.



**Figure 3. Results of LASSO regression predicting alcohol sipping after 12 months**


1   **A.** Rate of alcohol sipping after 12 months in each group. **B.** Estimates of coefficients of the survived  
2   variables (x axis: coefficient estimates; y axis: predictor). Site variables were excluded for a clarity  
3   (see **Figure S2** for the site variables). **C.** Distribution of the area under the curve (AUC) values (left)  
4   and a representative receiver-operating characteristic (ROC) curve (right) for the training and test  
5   datasets.

6   \*:  $p<0.05$ , \*\*:  $p<0.01$ ; \*\*\*:  $p<0.001$



**Figure 4.** Caffeinated soda consumption at baseline as a mediator of the relationship between impulsivity and alcohol sipping after 12 months

1 Along with family risk factors, caffeinated soda intake was a significant mediator between  
2 impulsivity and future alcohol sipping. Significant paths are flagged with asterisks (\*:  $p<0.05$ , \*\*:  
3  $p<0.01$ ; \*\*\*:  $p<0.001$ ). Indirect effects of caffeinated soda intake and family risk factors on the  
4 relationship between impulsivity and alcohol sipping after 12 months were significant (caffeinated  
5 soda intake:  $B=0.002$ , 95% CI=[0.001, 0.003],  $p=0.013$ ; family risk factors:  $B=-0.002$ , 95%  
6 CI=[-0.003, -0.001],  $p=0.006$ ).



1 **Table 1.** Results of hierarchical regression analysis predicting alcohol sipping after 12 months

Variables	Model 1	Model 2	Model 3
<b>Sex (Male)</b>	-0.020 (-0.031 - -0.010)***	-0.015 (-0.025 - -0.005)**	-0.013 (-0.023 - -0.003)**
<b>Race (Caucasian)</b>	0.032 (0.011 - 0.053)**	0.022 (0.002 - 0.042)*	0.022 (0.002 - 0.042)*
<b>Race (African American)</b>	0.001 (-0.024 - 0.026)	0.003 (-0.021 - 0.027)	0.003 (-0.021 - 0.027)
<b>Family income</b>	0.007 (0.000 - 0.015)*	0.005 (-0.002 - 0.012)	0.005 (-0.002 - 0.012)
<b>Parental education</b>	0.009 (0.002 - 0.015)*	0.007 (0.001 - 0.014)*	0.009 (0.002 - 0.015)*
<b>Parental monitoring</b>	-0.003 (-0.009 - 0.002)	-0.002 (-0.007 - 0.003)	-0.001 (-0.006 - 0.004)
<b>Family history of drug</b>	-0.003 (-0.015 - 0.009)	-0.005 (-0.017 - 0.007)	-0.005 (-0.017 - 0.006)
<b>Family history of alcohol</b>	-0.008 (-0.019 - 0.003)	-0.010 (-0.020 - 0.001)	-0.010 (-0.021 - 0.001)
<b>BMI</b>	0.007 (0.002 - 0.012)**	0.006 (0.001 - 0.011)*	0.006 (0.001 - 0.011)*
<b>Physical activity</b>	0.007 (0.001 - 0.012)*	0.005 (0.000 - 0.010)	0.005 (0.000 - 0.011)
<b>Lack of sleep</b>	0.000 (-0.005 - 0.006)	0.002 (-0.004 - 0.007)	0.001 (-0.004 - 0.007)
<b>Externalizing behaviors</b>	0.009 (0.003 - 0.015)**	0.007 (0.001 - 0.013)*	0.006 (0.000 - 0.013)*
<b>Internalizing behaviors</b>	-0.003 (-0.009 - 0.004)	-0.003 (-0.009 - 0.003)	-0.003 (-0.009 - 0.003)
<b>Alcohol sipping (Baseline)</b>		0.174 (0.161 - 0.187)***	0.172 (0.159 - 0.185)***
<b>Monthly caffeinated soda intake</b>			0.021 (0.010 - 0.032)***
<b>Weekly caffeinated soda intake</b>			0.024 (0.008 - 0.041)**
<b>Daily caffeinated soda intake</b>			0.036 (0.010 - 0.062)**

2 *Note.* The values are standardized regression coefficients (95% confidence intervals) from each  
3 step in the hierarchical analyses predicting alcohol sipping after 12 months. Significant variables  
4 predicting alcohol sipping are flagged with asterisks (\*: p<0.05, \*\*: p<0.01; \*\*\*: p<0.001). Higher  
5 consumption of caffeinated soda intake (Model 3) predicted a higher tendency for alcohol sipping  
6 after 12 months even after controlling for the confounding variables (13 variables from the top and  
7 21 data collection sites in Model 1) and alcohol sipping collected at baseline (Model 2).

8 *Abbreviations.* BMI, Body Mass Index.

9